

2024 HIV DRUG CHART

This quick-reference chart compares antiretroviral (ARV) options for the treatment of HIV, including adult dosing and dietary restrictions. Visit poz.com/drugchart for more info.

*Generic version available in the U.S. (Pills not shown actual size)

Complete Regimens



ATRIPLA*
efavirenz + tenofovir disoproxil fumarate + emtricitabine
One tablet once a day. Each tablet contains 600 mg efavirenz + 300 mg tenofovir disoproxil fumarate + 200 mg emtricitabine. Take on an empty stomach. Dose should be taken at bedtime to minimize dizziness, drowsiness and impaired concentration.



BIKTARVY
bictegravir + tenofovir alafenamide + emtricitabine
One tablet once a day. Each tablet contains 50 mg bictegravir + 25 mg tenofovir alafenamide + 200 mg emtricitabine. Take with or without food.



CABENUVA
cabotegravir + rilpivirine
A long-acting injectable regimen administered as two intramuscular injections every four weeks or eight weeks. A one-month lead-in period with Vocabria (cabotegravir) + Edurant (rilpivirine) pills is optional. Take with food.



COMPLERA
rilpivirine + tenofovir disoproxil fumarate + emtricitabine
One tablet once a day. Each tablet contains 25 mg rilpivirine + 300 mg tenofovir disoproxil fumarate + 200 mg emtricitabine. Take with a meal.



DELSTRIGO
doravirine + tenofovir disoproxil fumarate + lamivudine
One tablet once a day. Each tablet contains 100 mg doravirine + 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Take with or without food.



DOVATO
dolutegravir + lamivudine
One tablet once a day. Each tablet contains 50 mg dolutegravir + 300 mg lamivudine. Take with or without food.



GENVOYA
elvitegravir + cobicistat + tenofovir alafenamide + emtricitabine
One tablet once a day. Each tablet contains 150 mg elvitegravir + 150 mg cobicistat + 10 mg tenofovir alafenamide + 200 mg emtricitabine. Take with food.

Nucleoside/Nucleotide Reverse Transcriptase Inhibitors (NRTIs, or nukes)



CIMDUO
tenofovir disoproxil fumarate + lamivudine
One tablet once a day. Each tablet contains 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Take with or without food.



DESCOVI
tenofovir alafenamide + emtricitabine
One tablet once a day. Each tablet contains 25 mg tenofovir alafenamide + 200 mg emtricitabine. Take with or without food.



EMTRIVA*
emtricitabine (also known as FTC)
One 200 mg capsule once a day. Take with or without food.



EPIVIR*
lamivudine (also known as 3TC)
One 300 mg tablet once a day, or one 150 mg tablet twice a day. Take with or without food. Also approved for the treatment of hepatitis B virus but at a lower dose. People living with both viruses should use the HIV dose.



EPZICOM*
abacavir + lamivudine
One tablet once a day. Each tablet contains 600 mg abacavir + 300 mg lamivudine. Take with or without food. Should be used only by individuals who are HLA-B*5701 negative.



TEMIXYS
tenofovir disoproxil fumarate + lamivudine
One tablet once a day. Each tablet contains 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Take with or without food.



TRUVADA*
tenofovir disoproxil fumarate + emtricitabine
One tablet once a day. Each tablet contains 300 mg tenofovir disoproxil fumarate + 200 mg emtricitabine. Take with or without food.



VIREAD*
tenofovir disoproxil fumarate
One 300 mg tablet once a day. Take with or without food.



ZIAGEN*
abacavir
One 300 mg tablet twice a day, or two 300 mg tablets once a day. Take with or without food. Should be used only by individuals who are HLA-B*5701 negative.

Protease Inhibitors (PIs)



EVOTAZ
atazanavir + cobicistat
One tablet once a day. Each tablet contains 300 mg atazanavir + 150 mg cobicistat. Take with food.



KALETRA*
lopinavir + ritonavir
Two tablets twice a day, or four tablets once a day, depending on HIV drug resistance. Each tablet contains 200 mg lopinavir + 50 mg ritonavir. Take with or without food.



PREZCOBIX
darunavir + cobicistat
One tablet once a day. Each tablet contains 800 mg darunavir + 150 mg cobicistat. Take with food.



PREZISTA
darunavir
One 800 mg tablet, or two 400 mg tablets plus one 100 mg Norvir tablet once a day, or one 600 mg tablet plus one 100 mg Norvir tablet twice a day, depending on drug resistance. Take with food.



REYATAZ*
atazanavir
Two 200 mg capsules once a day, or one 300 mg capsule plus one 100 mg Norvir tablet once a day. Take with food.

Integrase Inhibitors



ISENTRESS
raltegravir
Two 600 mg Isentress HD tablets (shown) once a day for those who are treatment naive or whose virus has been suppressed on an initial regimen of Isentress. One 400 mg Isentress tablet twice daily for people with HIV treatment experience. Take with or without food.



TIVICAY
dolutegravir
One 50 mg tablet once a day for those first starting ARV therapy or for those who have not used an integrase inhibitor in the past. One 50 mg tablet twice a day for people with treatment experience who have HIV that is resistant to other integrase inhibitors and when taken with certain ARVs. Take with or without food.



VOCABRIA
cabotegravir
One 30 mg tablet taken once a day with once-daily Edurant for a month as an optional lead-in regimen before switching to Cabenuva injections or for short-term treatment. Take with food.



JULUCA
dolutegravir + rilpivirine
 One tablet once a day. Each tablet contains 50 mg dolutegravir + 25 mg rilpivirine. Take with a meal.



ODEFSEY
rilpivirine + tenofovir alafenamide + emtricitabine
 One tablet once a day. Each tablet contains 25 mg rilpivirine + 25 mg tenofovir alafenamide + 200 mg emtricitabine. Take with a meal.



STRIBILD
elvitegravir + cobicistat + tenofovir disoproxil fumarate + emtricitabine
 One tablet once a day. Each tablet contains 150 mg elvitegravir + 150 mg cobicistat + 300 mg tenofovir disoproxil fumarate + 200 mg emtricitabine. Take with food.



SYMFI AND SYMFI LO
efavirenz + tenofovir disoproxil fumarate + lamivudine
 One tablet of either Symfi or Symfi Lo once a day. Each tablet of Symfi contains 600 mg efavirenz + 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Each tablet of Symfi Lo (shown) contains 400 mg efavirenz + 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Take on an empty stomach. Dose should be taken at bedtime to minimize dizziness, drowsiness and impaired concentration.



SYMITUZA
darunavir + cobicistat + tenofovir alafenamide + emtricitabine
 One tablet once a day. Each tablet contains 800 mg darunavir + 150 mg cobicistat + 10 mg tenofovir alafenamide + 200 mg emtricitabine. Take with food.



TRIUMEQ
dolutegravir + abacavir + lamivudine
 One tablet once a day. Each tablet contains 50 mg dolutegravir + 600 mg abacavir + 300 mg lamivudine. Take with or without food. Should be used only by individuals who are HLA-B*5701 negative.

Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs, or non-nukes)



EDURANT
rilpivirine
 One 25 mg tablet once a day. Take with food.



INTELENCE
etravirine
 One 200 mg tablet twice a day. Take with food.



PIFELTRO
doravirine
 One 100 mg tablet once a day. Take with or without food.



SUSTIVA *
efavirenz
 One 600 mg tablet (shown) once a day, or three 200 mg capsules once a day. Take on an empty stomach or with a low-fat snack. Dose should be taken at bedtime to minimize dizziness, drowsiness and impaired concentration.



RUKOBIA
fostemsavir
 One 600 mg tablet twice a day for people with HIV treatment experience. Take with or without food.



SELZENTRY
maraviroc
 One 150 mg, 300 mg (shown) or 600 mg tablet twice a day, depending on other meds used, for people with HIV treatment experience. Take with or without food.



TROGARZO
ibalizumab
 A long-acting injectable administered intravenously as a single loading dose of 2,000 mg followed by a maintenance dose of 800 mg every two weeks for people with HIV treatment experience.

PK Boosters




NORVIR *
ritonavir
 Norvir is usually taken to boost the levels of other ARVs in the blood. Take with food.



TYBOST
cobicistat
 One 150 mg tablet once a day in combination with ARVs that require boosting. Used only to boost other drugs. Take with food.

Capsid Inhibitors



SUNLENCA
lenacapavir
 Sunlenca tablets are taken as a loading dose, with injections once every six months thereafter. Take with or without food.

These antiretroviral medications are rarely prescribed and no longer recommended:



APTIVUS
 tipranavir



COMBIVIR *
 zidovudine + lamivudine



CRIXIVAN
 indinavir



FUZEON
 enfuvirtide



INVIRASE
 saquinavir



LEXIVA
 fosamprenavir



RETROVIR *
 zidovudine (AZT)



TRIZIVIR
 abacavir + zidovudine + lamivudine



VIRACEPT
 nelfinavir



VIRAMUNE
 nevirapine



ZERIT
 stavudine (d4T)

Visit poz.com/drugchart-prevention for a list of ARV options to prevent HIV.

You can live well with HIV

It's important to start HIV treatment as soon as possible and stick with it. Remember to take your treatment as prescribed and stay engaged in care. Taking care of yourself is a great way to help you live well with HIV.



There is power in maintaining your sexual health


HIV doesn't have to stop you from being you. Discover helpful tips and support to keep living your authentic life.

Find more helpful information and resources at [HelpStopTheVirus.com](https://www.HelpStopTheVirus.com)

STOP THE VIRUS.



GILEAD and the GILEAD Logo are trademarks of Gilead Sciences, Inc. All other marks are the property of their respective owners.
© 2022 Gilead Sciences, Inc. All rights reserved. US-UNBC-0983 11/22



Treatment helps prevent the spread of HIV

If you're living with HIV, a major goal is to get your viral load to undetectable. This means that there is so little virus in the blood that a lab test can't measure it. **Current research shows that taking treatment as prescribed and getting to and staying undetectable prevents the transmission of HIV through sex. This is also known as U=U.**

Why U=U matters

U=U means undetectable=untransmittable. It helps destigmatize living with HIV, raises awareness that today's medications can be effective, and reminds people of the importance of continuing to take treatment as prescribed. Stay empowered to live a longer and healthier life.



Speak with your healthcare provider

It's important to be open and honest with your healthcare provider to find the right treatment for you and your routine. No questions are off limits when you meet with them.




Model portrayal

**Start HIV
treatment
ASAP**

Learn how to look after your health

If you are living with HIV, talk to your healthcare provider about treatment options.

Help Stop the Virus provides resources and information that can help you stay engaged in your health. Visit [HelpStopTheVirus.com](https://www.HelpStopTheVirus.com)



SEE INSIDE



STOP THE VIRUS.